


Managed Care Withhold Measures 2014-2015 - CHILDREN

| <b><u>Asthma Optimal Care</u></b>                                     |   |
|---|---|
| Medication Management for People with Asthma (MMA)                    | The percentage of patients 5-64 years of age during the measurement year who were identified as having persistent asthma and were dispensed appropriate medications that they remained on during the treatment period. Two rates are reported: (1.) The percentage of patients who remained on an asthma controller medication for at least 50% of their treatment period. (2.) Remained on an asthma controller medication for at least 75% of their treatment period.   |
| Use of Appropriate Medication for People with Asthma (ASM)            | The percentage of patients 5-64 years of age during the measurement year who were identified as having persistent asthma and who were appropriately prescribed medication during the measurement year.  |
| <b><u>Preventive Health Optimal Care</u></b>                          |   |
| Well-Child Visits in the First 15 Months of Life                      | The percentage of patients who turned 15 months old during the measurement year and who had the following number of well-child visits with a primary care physician during their first 15 months of life (6 visits).  |
| Well-Child Visits in the Third, Fourth, Fifth and Sixth Years of Life | The percentage of patients 3-6 years of age who had one or more well-child visits with a primary care physician during the measurement year.  |
| Adolescent Well-Care Visits   | The percentage of patients 12-21 years of age who had at least one comprehensive well-care visit with a primary care physician or an OB/GYN practitioner during the measurement year.   |
| <b><u>Behavioral Health Measures</u></b>                              |   |
| Follow-Up Care for Children Prescribed ADHD Medication (ADD)          | The percentage of children newly prescribed attention-deficit/hyperactivity disorder (ADHD) medication who had at least three follow-up care visits within a 10-month period, one of which was within 30 days of when the first ADHD medication was dispensed. The reported rate, the Initiation Phase, is the percentage of members 6-12 years of age as of the IPSP with an ambulatory prescription dispensed for ADHD medication, who had one follow-up visit with practitioner with prescribing authority during the 30-day Initiation Phase. |



# Coding of HEDIS® Well-Child and Adolescent Visits




Documentation should include:

- » Physical examination
- » Growth and development assessment
- » Age-Appropriate anticipatory guidance (can include physical activity and nutrition)

This flowchart illustrates the components of a HEDIS® well visit. Improving coding and documentation are two ways to improve your HEDIS® scores.

Use of preventive (993XX) procedure code **AND/OR** diagnosis code listed meets well visit coding requirement. Maximize visits by adding appropriate immunizations, screenings and age-appropriate anticipatory guidance.

# The Flow of HEDIS® Well-Child and Adolescent Visits



This flowchart illustrates the components of a HEDIS® well visit. Improving coding and documentation are two ways to improve your HEDIS® scores.

Documentation should include:

- » Physical examination
- » Growth and development assessment
- » Age-Appropriate anticipatory guidance (can include physical activity and nutrition)